

POSTER PRESENTATION

Open Access

Peroneus brevis rupture associated with a hypertrophic peroneal tubercle: a case report and literature review

Dean Samaras*, Andrew Kingsford

From Australasian Podiatry Council Conference 2013 Sydney, Australia. 2-5 June 2013

Varying morphology of the peroneal tubercle can be associated with peroneal tendon pathology including tendonosis, stenosing tenosynovitis, tendon attrition and tears. The authors present a case of peroneus brevis rupture associated with an enlarged peroneal tubercle in a 59yo male. The clinical testing, medical imaging, diagnosis, surgical reconstruction and rehabilitation is discussed. The literature review results are also reported.

A literature review was conducted across several scientific databases including Medline, CINAHL and PubMed using the search terms "peroneus brevis" AND "rupture" AND "peroneal tubercle" OR "peroneal trochlea". Further information was sought with citation tracking and reference checking and reviewing unpublished data.

No report of isolated peroneus brevis rupture associated with a hypertrophic peroneal tubercle was found in the literature. However, there were several reports of peroneal tendinopathy associated with this pathology. Failing conservative care, good to excellent results were reported with surgical resection and tendon repair. In our case resection of the tubercle and peroneus brevisrepair with a split thickness peroneus longus graft was performed. At 3 years post-op the patient was asymptomatic, had returned to previous level of activity and was satisfied with the result.

The hypertrophic peroneal tubercle can cause significant lateral foot pain. It is important for clinicians to keep this pathologic process in the list of differential diagnoses. X-rays (particularly the axial calcaneal view), ultrasound, CT and/or MRI can identify the extent of the pathology. If conservative measures fail, generous

resection and repair of any peroneal tendinopathy can result in return to normal function.

Published: 31 May 2013

doi:10.1186/1757-1146-6-S1-P14

Cite this article as: Samaras and Kingsford: Peroneus brevis rupture associated with a hypertrophic peroneal tubercle: a case report and literature review. Journal of Foot and Ankle Research 2013 6(Suppl 1):P14.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit



^{*} Correspondence: deansamaras@gmail.com Kingsford Podiatry Group, Melbourne, Victoria, 3000, Australia

