### **ORAL PRESENTATION**



**Open Access** 

# Epidemiology of foot complaints in systemic lupus erythematosus

S Otter<sup>1\*</sup>, S Kumar<sup>2</sup>, P Gow<sup>2</sup>, N Dalbeth<sup>3</sup>, KA Davies<sup>4</sup>, S Panthakalam<sup>5</sup>, K Rome<sup>1</sup>

*From* Australasian Podiatry Conference 2015 Queensland, Australia. 6-8 May 2015

#### Background

Foot complaints are common in inflammatory arthropathies such as rheumatoid arthritis and cause considerable disability. However, little is known about the nature and extent of foot complaints in systemic lupus erythematosus (SLE) as highlighted by a recent systematic review [1]. We set out to explore the clinical features and symptoms of foot involvement among people with SLE from the patients' perspective.

#### Methods

We developed a new 40-item item self-administered questionnaire, posted to patients with SLE attending adult rheumatology clinics at Auckland and Counties Manukau District Health Boards, Auckland, New Zealand. The questionnaire enquired about symptoms of foot pain, extra-articular features, assessed the anatomical distribution of symptoms according to validated foot-mannequins and considered the impact of symptoms on subjects activities of daily living and well-being.

#### Results

A total of 107 responses were received, with a mean age of 52 years (SD 14.1) and a mean duration of positive diagnosis of SLE of 12 years (SD10.9). Overall 79% of respondents reported foot pain caused by SLE with 50% reporting current foot pain. All regions of the feet were affected, with the midfoot, hindfoot and ankles most troublesome. Other symptoms including swelling, numbness. soft tissue pain and cold feet were common. For 38% foot pain stopped them sleeping and in 35% foot pain negatively affected them emotionally; overall for 64% foot pain generally adversely affected their lives.

\* Correspondence: simon.otter@aut.ac.nz

<sup>1</sup>Auckland University of Technology, Auckland, New Zealand Full list of author information is available at the end of the article Only 27% reported social and/or family activities were never impacted by foot complaints.

#### Conclusions

For the first time we have characterised foot complaints in a large group of people with SLE. Foot symptoms in SLE are common, severe and heterogeneous in nature, typically having a considerable negative impact on patient's well-being.

#### Authors' details

<sup>1</sup>Auckland University of Technology, Auckland, New Zealand. <sup>2</sup>Middlemore Hospital, Papatoetoe 2025, New Zealand. <sup>3</sup>The University of Auckland, Auckland, 1010, New Zealand. <sup>4</sup>Brighton & Sussex Medical School, Brighton, BN1 9PX, UK. <sup>5</sup>Eastbourne District General Hospital, Eastbourne, BN21 2UD, UK.

#### Published: 22 September 2015

#### Reference

 Williams A, Grofts G, Teh L: 'Focus on feet' the effects of Systemic Lupus Erythematosus - a narrative view of the literature. Lupus 2013, 22:1017-23.

#### doi:10.1186/1757-1146-8-S2-O30

**Cite this article as:** Otter *et al.*: **Epidemiology of foot complaints in systemic lupus erythematosus.** *Journal of Foot and Ankle Research* 2015 **8** (Suppl 2):O30.

## Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

) BioMed Central

Submit your manuscript at www.biomedcentral.com/submit



© 2015 Otter et al. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http:// creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/ zero/1.0/) applies to the data made available in this article, unless otherwise stated.